

Harry's Beau&Bon Semi-complet

Environmental

Product Declaration



The first EPD process certified in the Food industries







REGISTRATION NUMBER

S-P-01561

CPC CODE234 BAKERY
PRODUCTS
PCR 2012:06 VER. 3.0
20/01/2020

PUBLICATION DATE

2020/07/30

REVISION

2 of 2021/12/22

VALID UNTIL

2025/07/12

PROGRAMME

The International EPD® System www.environdec.com

PROGRAMME OPERATOR

EPD International AB

This EPD has been developed in conformity to ISO 14025. An EPD should provide current information and may be updated if conditions change. The stated validity is, therefore, subject to the continued registration and publication at www.environdec.com.



1. Brand and product

THE BRAND HARRYS

Launched in 1970 on the baked goods market in France, the Harrys brand today is leader in the soft bread market and a important player in the morning goods market. Quality and innovation are two of the most important key success factors.

PLANT AND PROCESS

Beau&Bon Semi-complet is produced in the French plant of Valenciennes, where a typical bakery process takes place.


The process steps involved in the manufacture are: mixing of the ingredients, to form a dough which will be divided and rounded, dough molding and proofing, pieces baking in a specific oven, cooling and cutting in slices.

Beau&Bon Semi-complet is packed into 400 grams format and it is ready for consumption.

More info on www.harrys.fr



THE PRODUCT



NUTRITIONAL INFORMATION (per 100 g)							
Energy	kJ kcal	1 157 274					
Fats of which saturated	grams	3.9 0.4					
Carbohydrates of which sugars	grams	48.9 6.2					
Fibres	grams	5.7					
Proteins	grams	8					
Salt	grams	1					






2. Barilla group

Passion for quality, continuous pursuit of excellent recipes and ability to combine tradition and innovation are the fundamental ingredients that that have allowed a small shop of bread and pasta, opened in 1877 in Parma, to become an international player in the market of pasta, ready-to-eat sauces, baked goods and crispy breads.

The Group operates in over 100 countries through its brands, which have become the icon of excellence in the food sector, and with 30 production sites, which every year contribute to the production of over 2,099,000 tonnes of products.

With its brands - Barilla, Mulino Bianco, Pan di Stelle, Gran Cereale, Harrys, Pavesi, Wasa, Filiz, Yemina e Vesta, Misko, Voiello, Cucina Barilla, Catelli, Lancia, Tolerant and Pasta Evangelists – promotes a tasty, joyful and healthy diet, inspired by the Mediterranean diet and the Italian lifestyle.

Further information on www.barillagroup.com



Good for You, Good for the Planet

In order to make a concrete contribution to global challenges, over the years, Barilla has developed a thought enclosed in the Good for You, Good for the Planet Mission that guides, step by step and offers people good, safe, nutritionally balanced food, coming from responsible supply chains.

GOOD FOOD means taste, pleasure and a daily gesture of love for the people themselves.

HEALTHY FOOD means selected raw materials and balanced nutritional profiles to support healthy lifestyles.

FOOD SOURCED FROM RESPONSIBLE SUPPLY CHAINS means seeking the best ingredients to guarantee excellent quality, respectful of people, animals and the environment.

A commitment "from field to fork", which has led to the development of initiatives in the various stages of the supply chain and for which all Barilla Group brands contribute through projects aiming to improve the nutritional profile of products, reinforce the sustainability of the production and supply chains and provide transparent communication to consumers.

















































Harrys: a CO₂ Compensated Brand

Harrys brand has embarked on a path that has led to the achievement of **100% carbon compensation**, in line with its core values - quality, unique softness and local and responsible engagement.

The path includes three steps: **measure. reduce and compensate**. The path was certified by DNV-GL following the International Standard PAS 2060.

This certification covers both Harrys brand and products.





For more information, please visit: www.harrys.fr/moelleux-responsable/co2-compense



WE MEASURE

Starting from 2019 all the greenhouse gas emissions arising from Harrys brand activities (from field to shelf) are **yearly identified and measured**, related to a baseline year (2018).



WE REDUCE

Some measures are taken to reduce the greenhouse gases emissions, like Energy Saving Programs, Charte Harrys project for a more sustainable soft wheat farming and purchasing **renewable electricity** (100% from hydropower sources).



WE COMPENSATE

To compensate the remaining emissions, Harrys has chosen to contribute to the protection of a rainforest and support solar energy use through projects labelled by the **Verified Carbon Standard** (VCS) and the **Climate Community and Biodiversity Alliance Standard** (CCBA). Projects developed under these programs must follow a rigorous assessment process in order to be certified. Harrys has chosen to contribute to the protection of rainforest in Brasil and support solar energy in India.







Harrys CO₂ compensation project

100%



Floresta de Portel

Brazilian forest is one of the richest ecosystems in the world and is now threatened.

The Floresta de Portel projects aims at protecting and preserving this fragile environment.



Solar India

The aim of this project is to supply renewable solar energy in India, by installing photovoltaic panels in Telangana and Maharashtra states.







3. Environmental performance calculation



The Environmental performance of the product was calculated using the LCA (life cycle analysis) methodology, including the entire production chain, starting from the cultivation of the raw materials until the delivery of the finished product to the shelf.

The study was conducted following the specific product rules published for the EPD System: "CPC code 234 – Bakery products".

The contribution to the environmental impacts brought by generic data is less than 10% in all impact categories.

DECLARED UNIT

Data are referred to 1 kg of product plus the related packaging (the packaging is referred to the 400 grams format, reported to 1 kg of product).

SYSTEM BOUNDARIES

The processes constituting the analyzed system were organized according to following three successive phases, in compliance with the EPD system's requirements.







4. Raw materials production



SOFT WHEAT FLOUR AND BRAN AND OTHER CEREALS

Cereal cultivation performances are calculated on the basis of primary data, yield, energy consumption and fertiliser use for each cereal, collected from farms. Cultivation region is France.

Cultivation yield is calculated as average of years 2018, 2019 and 2020.

VEGETABLE OILS

Data for sunflower oil cultivation come from secondary data (collected from Agrifootprint database), the extraction and refinery data come from literature (Nilsson et al. 2010).

Data for rapeseed oil cultivation come from secondary data (collected from Agrifootprint database) and the refinery data come from literature (Schmidt 2007).



INGREDIENTS PRODUCTION

OTHER INGREDIENTS

Data related to sugar production are primary and come from Barilla suppliers.

Data from LCA databases are used for other material in the recipe (yeast, salt and vinegar).





5. Packaging and auxiliary materials production




PACKAGING PRODUCTION

PRIMARY PACKAGING

Packaging environmental performances are calculated using the 400 g format, the only format for the product, and reported per packaging used for 1 kg of product.

The primary packaging consists in a plastic film sac with a plastic based closure clip.

Primary data (from packaging unit) are used for packaging amount and packaging materials production; data about packaging production process come from Barilla LCA database.



LCA Pack Designer

Since 2004, Barilla designs new packaging with esign tool". It allows the environmental

the "LCA packaging design tool". It allows the assessment of the environmental impacts of the packaging solutions already during the design phase.

PACKAGING FOR TRANSPORT

The packaging for transport consists in cardboard boxes (american box), used for the distribution of the product, and a plastic extensible film. Boxes are made mainly by recycled cardboard carton (pre and post consumer).

The data used have been collected by LCA database (mainly Ecoinvent).

Packaging used for Harrys products is 100% designed for recycle.

Auxiliary materials environmental performances are evaluated by using primary data from plant, during 2020 year. Secondary data (Ecoinvent) are used for environmental aspects associated to materials production.





6. Beau&Bon Semi-complet production



GENERAL INFORMATION

The environmental performances related to the production process are evaluated considering as primary data the energy and the water consumption and the waste production. Secondary data (mainly Ecoinvent) are used for the environmental aspects related to the production of energy and water.

The plant considered in the analysis is Valenciennes.

WATER

The water consumption is evaluated using primary data. The overall value is attributed to the product using the mass allocation procedure.


Plant water consumption includes also the water amount needed for dough preparation: this amount is included both in plant consumption and product recipe following a precautionary approach.

Data are referred to year 2020

WASTE

The primary data are collected by the plant registrations. The overall value is attributed to the product using the mass allocation procedure.

Data are referred to year 2020.



AMERICAN SANDWICH NATURE PRODUCTION

ELECTRICITY

Total plants electricity has been divided using mass allocation (plants produce other products beyond Beau&Bon Semi-complet).

Barilla, through the GO certification system (Guaranty of origin market), buys energy from hydroelectric renewable resources as to cover the entire Harrys production. Data are referred to 2020.

NATURAL GAS

The natural gas consumption is evaluated using primary data. The overall value is attributed to the product using the mass allocation procedure.

Data are referred to year 2020.





7. Distribution



DISTRIBUTION

Beau&Bon Semi-complet is produced in the French plant of Valenciennes and it is sold mainly in France.

Distribution performance were calculated considering the transport in France for about 807 km.

All transport stage from plant to retailer are included.

The product does not need any particular storage condition (such as refrigeration).


The impacts related to the disposal of the packaging for transport have been calculated considering the scenario for paperboard and plastic for distribution in France (reference: Eurostat 2018).







8. Packaging end of life



* Data elaborated from Eurostat database, reference year 2018





9. Environmental results

USE OF RESOURCES data referred to 1 kg of product		UPSTREAM		CORE	DOWNSTREAM		
		Raw material production	Packaging and auxiliary materials production	Production	Distribution up to shelf	Primary packaging end of life	TOTAL
PRIMARY ENERGY RESOURCES - RENEWABLE data in MJ	Used as energy carrier	7,37E-01	5,24E-01	3,96E+00	2,74E-03	2,54E-05	5,23E+00
	Used as raw materials*	0,00E+00	2,20E-01	0,00E+00	0,00E+00	0,00E+00	2,20E-01
	Total	7,37E-01	7,44E-01	3,96E+00	2,74E-03	2,54E-05	5,45E+00
PRIMARY ENERGY RESOURCES - NON RENEWABLE	Used as energy carrier	5,13E+00	3,48E+00	4,42E+00	1,77E+00	1,54E-03	1,48E+01
	Used as raw materials	8,02E-05	5,78E-01	0,00E+00	0,00E+00	0,00E+00	5,78E-01
data in MJ	Total	5,13E+00	4,06E+00	4,42E+00	1,77E+00	1,54E-03	1,54E+01
Seconda	Secondary Material (g)		1,24E+02	0,00E+00	0,00E+00	0,00E+00	1,24E+02
Renewable secondary fuels (MJ. net calorific power)		0,00E+00	7,38E-02	0,00E+00	0,00E+00	0,00E+00	7,38E-02
Non-renewable secondary fuels (MJ. net calorific power)		0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
Net use of fresh water (liters)		2,21E+01	1,44E+00	3,15E+01	4,03E-02	6,28E-04	5,50E+01
OUTPUT FLOWS data referred to 1 kg of product		UPSTREAM		CORE	DOWNSTREAM		
		Raw material production	Packaging and auxiliary materials production	Production	Distribution up to shelf	Primary packaging end of life	TOTAL
Waste to anim	Waste to animal feed or similar (g)		0,00E+00	7,23E+02	0,00E+00	0,00E+00	7,23E+02
Components for reuse (g)		0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
Materials for recycling (g)		1,16E+00	1,68E+01	3,46E+01	1,07E+02	5,77E+00	1,65E+02
Materials for energy recovery (g)		2,24E-02	0,00E+00	0,00E+00	5,11E+00	8,03E+00	1,32E+01
Exported energy. electricity (MJ)		0,00E+00	0,00E+00	0,00E+00	2,20E-03	4,40E-03	6,60E-03
Exported energy. thermal (MJ)		0,00E+00	0,00E+00	0,00E+00	4,60E-03	9,20E-03	1,38E-02
Secondary energy resources and recovered energy flows do not show relevant contributions. *The biomasses transformed into the product are not considerable product are not considerable product are not considerable product.							uct are not considered.





POTENTIAL ENVIRONMENTAL IMPACTS data referred to 1 kg of product		UPSTREAM		CORE	DOWNSTREAM		
		Raw material production	Packaging and auxiliary materials production	Production	Distribution up to shelf	Primary packaging end of life	TOTAL
GLOBAL WARMING POTENTIAL - GWP (g CO ₂ eq)	Fossil	4,78E+02	1,72E+02	2,90E+02	1,27E+02	1,42E+01	1,08E+03
	Biogenic	5,14E+00	2,61E-01	2,13E+00	4,57E+00	4,71E-02	1,21E+01
	Land use and land transformation	4,05E+01	1,85E+00	1,93E-03	1,11E-03	2,54E-05	4,24E+01
	Total	5,24E+02	1,74E+02	2,92E+02	1,32E+02	1,43E+01	1,14E+03
Acidification Potentia	al - g SO ₂ eq.	7,41E+00	6,24E-01	6,31E-01	3,88E-01	1,80E-03	9,05E+00
Eutrophication Poter	ntial - g PO ₄ eq.	4,89E+00	1,68E-01	9,25E-02	5,24E-02	5,35E-04	5,20E+00
Photochemical Oxida	ant Formation Potential - gNMVOC eq	1,45E+00	6,05E-01	7,55E-01	4,29E-01	2,54E-03	3,24E+00
Abiotic Depletion Pot	tential - Elements g Sb eq.	7,93E-04	3,09E-05	5,79E-06	5,47E-06	4,86E-08	8,35E-04
Abiotic Depletion Pot value	tential - Fossil fuels - MJ, net calorific	4,24E+00	3,66E+00	4,38E+00	1,77E+00	1,51E-03	1,41E+01
Water scarcity poten	tial, m3 eq.	2,53E-01	4,93E-02	2,27E-01	-2,12E-04	1,97E-05	5,29E-01
		UPSTREAM		CORE	DOWNSTREAM		
111	STE PRODUCTION* Ferred to 1 kg of product	Raw material production	Packaging and auxiliary materials production	Production	Distribution up to shelf	Primary packaging end of life	TOTAL
Hazard	lous waste disposed (g)	7,19E-03	1,86E+00	0,00E+00	0,00E+00	0,00E+00	1,9E+00
Non-Haza	ardous waste disposed (g)	3,93E+00	2,03E+01	0,00E+00	0,00E+00	0,00E+00	2,4E+01
Radioa	ctive waste disposed (g)	9,49E-01	4,25E-01	9,96E-02	5,76E-02	4,42E-05	1,5E+00

The biogenic contribution to Global Warming Potential refers only to biogenic methane.


The contribution given by biogenic CO_2 is equal to zero, since the absorbed amount is equal to the emitted biogenic CO_2 within the reference 100 years period.

*Non-zero values indicate waste flows to disposal whose treatment impact isn't evaluated within system boundaries (usually they come from secondary data used in calculation model). Zero values indicate that – even if some waste are produced and disposed – their impact is evaluated within the system boundaries.





PRODUCT ENVIRONMENTAL PERFORMANCES



Compared to the last EPD, in this section the Water Scarcity indicator has substituted the Virtual Water Content, previously reported, to improve coherence with the indicators section.





10. Differeces versus previous versions of EPD

The differences versus previous EPD version are due mainly to the use of updated emission factors for the energy mixes, updated yields for soft wheat cultivation calculated as average value of the last three available years for every region. Moreover, the product Environmental performances

section has been modified with the substitution of Virtual Water Content with Water Scarcity indicator.

The product name has changed: in 2020 it was "Harry's Pain de Mie", in 2021 the same product is sold as "Beau&Bon Semi-complet".

11. Additional information

REFERENCES

- International EPD Consortium, General Programme Instructions (EPD), ver. 3.01 of 18/09/2019;
- WWF, Global Footprint Network, Zoological Society of London, Living Planet Report 2008, WWF (2008);
- PCR 2012:06 CPC 234: Bakery Products; ver. 3.0 of 20/01/2020;
- Nilsson K., Flysjö A., Davis J., Sim S., Unger N., Bell S. "Comparative life cycle assessment of margarine and butter consumed in the UK, Germany and France" 2010, Int J Life Cycle Ass vol. 15 num. 9 p 916-926;
- Schmidt J.H. Life Cycle Assessment of rapeseed oil and palm oil 2010, International Journal of LCA 15 pp.183-197;
- Eurostat database for waste management, latest version (2018).



Environmental declarations published within the same product category, though originating from different programs. may not be comparable. This declaration and further information in regards are available at www.environdec.com





As EPD owner, Barilla has the sole ownership, liability and responsibility for the EPD.

EPD PROCESS CERTIFICATION

Product category Rules (PCR) review conducted by: Technical Committee of the International EPD® system. Chair Filippo Sessa

Contact via info@environdec.com

Program operator:

EPD International AB

Box 210 60, SE-100 31 Stockholm, Sweden

info@environdec.com



EPD PROCESS CERTIFICATION

Independent verification of the declaration and data, according to ISO 14025:



EPD process verification

EPD verification- Third party verifier

PROCESS INTERNAL VERIFICATION

Procedure for follow-up of data during EPD validity involves third part verifier:



Yes



No

Third party verifier: Bureau Veritas Certification Sweden AB, Accredited by: SWEDAC



Process internal verifier: Ugo Pretato, Approved by: The International EPD® System



CONTACTS

Barilla G. e R. Fratelli- Società per Azioni, via Mantova 166, 43122, Parma, Italy. www.barillagroup.com
For additional information relative to the activities of the Barilla Group or in regards to this environmental declaration, please contact:
Laura Marchelli - laura.marchelli@barilla.com



Technical support and grafic design: Life Cycle Engineering SpA - Italy www.lcengineering.eu







12. Glossary

ECOLOGICAL FOOTPRINT

The ecological footprint measures the area of biologically productive land and water required to provide the resources used and absorb the carbon dioxide waste generated along the entire life cycle. It is measured in standard units called global hectares (gha).

www.globalfootprint.org

ACIDIFICATION (AP)

It is a phenomenon for which precipitation is unusually acidic, meaning that it has substandard levels of pH. It can have harmful effects on plants, aquatic animals and infrastructure. Acid rain is caused by emissions of SO_2 . NO_x and NH_3 . The acidification potential is measured in mass of sulphur dioxide equivalent (SO2-eq).

CARBON FOOTPRINT

A product carbon footprint is the total amount of green-house gases produced along the entire life cycle. It is expressed in equivalent mass of carbon dioxide (CO₂-eq). In agriculture a significant contribution is given by the emission of nitrous oxide (N2O) due to the fertilizers use. It is also known as Global Warming Potential (GWP).

www.ipcc.ch

EUTROPHICATION (EP)

It is an abnormal proliferation of vegetation in the aquatic ecosystems caused by the addition of nutrients into rivers. lakes or ocean. which determinates a lack of oxygen. The eutrophication potential is mainly influenced by emission into water of phosphates and nitrates. It is expressed in mass of PO, "equivalent.

WATER SCARCITY

Water scarcity measures the available water remaining per unit of surface in a given watershed relative to the world average, after human and aquatic ecosystem demands have been met. This method builds on the assumption that the potential to deprive another user of water is directly proportional to the amount of water consumed and inversely proportional to the available water remaining per unit of surface and time in a region (watershed).

www.wulca-waterlca.org

PHOTOCHEMICAL OXIDANT FORMATION POTENTIAL (POFP)

Production of compounds that, under the light effect, are able to promote an oxidation reaction leading to ozone production in the troposphere.

The indicator is mainly influenced by VOCs (Volatile organic compounds) is usually expressed in mass of VOCs equivalent (g NMVOC - equivalent).

